Express Date of D	Deposit	:: July		7502US	01P	E SCHOOL STREET		ſ	Page 1 o	f 1
rease type	i i prius .			work Reduction Act	<b>.</b>	Patent and Tra	Approved for use t demark Office - U.S. D formation unless it disp	LPAR IMEN	FOE COMMI	RCF
Modified F	orm 144	9:PTO				Application Number	09/546,2	69	RE	CF
						Filing Date	04/10/00	ı		
	INFO	RMAT	ION DISC	LOSURE		First Named Inventor	Hunzikei			
	STA	TEMEN	IT BY API	PLICANT		Group Art Unit	1651		TEO: 6	
						Examiner Name	Witz, J.		TECH	ÆNTE:
	(use	as many	sheets as n	ecessary)		Attorney Docket Number	17811-0	14 CIP (	M-014 CI	P)
				· · · · · · · · · · · · · · · · · · ·	11.6	PATENT DOCUMENTS				
Exam	Cite	U.S. P		Issue Date	Γ	of Patentee(s) or Applicant(s)	Class	Sub	Filing D	ate
Initials	No.		ment No.	<u> </u>			Class	Class	If Appro	priate
74	A2	US 5,	968,546	10/19/99	Baur,	Marcus			05/15/9	8
					FORE	GN PATENT DOCUMENTS				
Exam Initials	Cite No.	Foreig Office	n Patent D Number	ocument	Name o	of Applicant(s)	Date of Publication	on	Trans Yes	lation No
$\rightarrow \mathcal{K}$	B1 ·	DE	196 51 99	92 A1	Tolocz	yki, Christian	06/25/98			Х
4	B2	wo	93/08776		Culture	Technology	05/13/93			
			0	THER PRIOR	ART -	NON PATENT LITERATURE DOCI	IMENTS			
Exam, Initials	Cite No.	Name				te), Publication, Volume, Page(s), D				
	C37					<u> </u>				
- W.	C38	International Search Report for International application No. PCT/IB00/01076  Lenoir-Viale, M.C. et al., 1993. "Epidermis reconstructed from the outer root sheath of human hair follicle: Effect of retinoic acid." Archives of Dermatological Research, vol. 285, no. 4, pages 197-204.						ffect of		
Ty.	C39	Limat, Alain et al., 1996. "Successful treatment of chronic leg ulcers with epidermal equivalents generated from cultured autologous outer root sheath cells." J. of Inv. Derm., vol. 107, no. 1, pages 128-135.						rom		
		]								


а сор, с	time reference to met promaca ac i	was providedly sited by or submitted to the silico in a prior application,
Serial No.	, filed	, and relied upon for an earlier filing date under
35 U.S.C.	§120 (continuation, continuation-in	-part, and divisional applications).

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered.

Include copy of this form with next communication to applicant.

TRA 1544872v1

## Express Mail No. EK674915 US Date of Deposit: July 10, 2000



Application Number	09 546.269
Filing Date	April 10, 2000
Applicants	Hunziker, et al
Group Art Unit	1636
Examiner Name	Not Yet Assigned
Attorney Docket Number	17811-014 (M-14 CIP)

Evam Initials	Cite No.	U.S. Patent Document No.	Issue Date	Name of Patentees	Class	Sub Class	Filing Date If Appropriate
AV	Αl	*5,580,781	12 3 1996	Naughton et al.	435	1.)	
				FOREIGN PATENT DOCUMEN	rs -		
Exam	Cite No.	Foreign Patent l	Document	Name of Applicants	Date of Pi	ıblication	Translation Yes No

Same of Author, Fille (when appropriate), Publication, Volume, Page(s), Date, Etc.	Exam	Cite	OTHER NON PATENT LITERATURE DOCUMENTS
C2 *Fabre, 29 Immunol, Lett. 161-166 (1991)  C3 *Harris et al., 18 Clin. Exp. Dermatol, 417-420 (1993)  C4 *Hetton et al., 14 J. Am. Acad. Dermatol, 399-405 (1986)  C5 *Hunyadi et al., 14 J. Dermatol, Surg. Oncol, 75-78 (1988)  C6 *Johnson et al., 11 J. Burn Care Rehab, 504-509 (1990)  C7 *Leigh et al., 117 Brit, J. Dermatol, 591-597 (1987)  C8 *Leigh et al., 11 Clin. Exp. Dermatol, 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol, 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol, 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258, (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Folliele: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)		No.	Name of Author, Title (when appropriate), Publication, Volume, Page(s), Date, Etc.
C3 *Harris et al., 18 Clin. Exp. Dermatol. 417-420 (1993)  C4 *Hetton et al., 14 J. Am. Acad. Dermatol. 399-405 (1986)  C5 *Hunyadi et al., 14 J. Dermatol. Surg. Oncol. 75-78 (1988)  C6 *Johnson et al., 11 J. Burn Care Rehab. 504-509 (1990)  C7 *Leigh et al., 11 Clin. Exp. Dermatol. 591-597 (1987)  C8 *Leigh et al., 11 Clin. Exp. Dermatol. 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology. 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology. 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts. Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular Ketatining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells. Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.	7 V		*Brysk <i>et al.</i> , 25 J. Am. Acad. Dermatol. 238-244 (1991)
C4 *Hetton et al., 14 J. Am. Acad. Dermatol. 399-405 (1986)  C5 *Hunyadi et al., 14 J. Dermatol. Surg. Oncol. 75-78 (1988)  C6 *Johnson et al., 11 J. Burn Care Rehab. 504-509 (1990)  C7 *Leigh et al., 117 Brit. J. Dermatol. 591-597 (1987)  C8 *Leigh et al., 11 Clin. Exp. Dermatol. 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY, The Journal of Investigative Dermatology. 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation, The Journal of Cell Biology. 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C18 Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C2	*Fabre, 29 Immunol. Lett. 161-166 (1991)
C5 *Hunyadi et al., 14 J. Dermatol. Surg. Oncol. 75-78 (1988)  C6 *Johnson et al., 11 J. Burn Care Rehab. 504-509 (1990)  C7 *Leigh et al., 117 Brit. J. Dermatol. 591-597 (1987)  C8 *Leigh et al., 11 Clin. Exp. Dermatol. 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PHLOSEBACEOUS APPARATUS TO CONTROLLED INJURY, The Journal of Investigative Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation, The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat et al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)  C16 Limat et al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.	4	C3	*Harris et al., 18 Clin. Exp. Dermatol. 417-420 (1993)
C6 *Johnson et al., 11 J. Burn Care Rehab. 504-509 (1990)  C7 *Leigh et al., 11 T Brit. J. Dermatol. 591-597 (1987)  C8 *Leigh et al., 11 Clin. Exp. Dermatol. 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts. Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C4	*Hetton <i>et al.</i> , 14 J. Am. Acad. Dermatol. 399-405 (1986)
C7 *Leigh et al., 117 Brit. J. Dermatol. 591-597 (1987)  C8 *Leigh et al., 11 Clin. Exp. Dermatol. 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells. Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C5	*Hunyadi et al., 14 J. Dermatol. Surg. Oncol. 75-78 (1988)
*Leigh et al., 11 Clin, Exp. Dermatol, 650-652 (1986)  C9 *Mol et al., 24 J. Am. Acad. Dermatol, 77-82 (1991)  C10 *Moll et al., 46 Hautarzt 548-552 (1995)  C11 *Phillips et al., 21 J. Am. Acad. Dermatol, 191-199 (1989)  C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY, The Journal of Investigative Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258, (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation, The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat et al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194; 218-227 (1991)  C16 Limat et al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C6	*Johnson et al., 11 J. Burn Care Rehab. 504-509 (1990)
<ul> <li>*Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)</li> <li>*Moll et al., 46 Hautarzt 548-552 (1995)</li> <li>*Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)</li> <li>Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology, 15:145-155 (1955)</li> <li>Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197</li> <li>C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)</li> <li>C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)</li> <li>C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell &amp; Tissue Research, 275:169-176 (1994)</li> <li>Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)</li> <li>Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.</li> </ul>		C7	*Leigh <i>et al.</i> , 117 Brit. J. Dermatol. 591-597 (1987)
<ul> <li>*Moll et al., 46 Hautarzt 548-552 (1995)</li> <li>C11 *Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)</li> <li>C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology. 15:145-155 (1955)</li> <li>C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)</li> <li>C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology. 109:2295-2312 (1989)</li> <li>C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research. 194: 218-227 (1991)</li> <li>C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell &amp; Tissue Research. 275:169-176 (1994)</li> <li>C05 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells. Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)</li> <li>C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.</li> </ul>	+	C8	*Leigh <i>et al.</i> , 11 Clin, Exp. Dermatol, 650-652 (1986)
<ul> <li>*Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)</li> <li>*C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY, The Journal of Investigative Dermatology, 15:145-155 (1955)</li> <li>*C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)</li> <li>*C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)</li> <li>*C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)</li> <li>*C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell &amp; Tissue Research, 275:169-176 (1994)</li> <li>*C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis. Cell, 61:1329-1337 (1990)</li> <li>*C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.</li> </ul>		<u>C</u> 9	*Mol et al., 24 J. Am. Acad. Dermatol. 77-82 (1991)
C12 Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY, The Journal of Investigative Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation, The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.		CTO	*Moll <i>et al.</i> , 46 Hautarzt 548-552 (1995)
PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY. The Journal of Investigative Dermatology. 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts. Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.		СП	*Phillips et al., 21 J. Am. Acad. Dermatol. 191-199 (1989)
Dermatology, 15:145-155 (1955)  C13 Montagna et al., The Structure and Function of Skin 172-258, (Academic Press New York, NY 197)  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. The Journal of Cell Biology, 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts. Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of Keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C12	Eosem et al., RESPONSES OF THE SUPERFICIAL PORTION OF THE HUMAN
C13 Montagna et al., The Structure and Function of Skin 172-258. (Academic Press New York, NY 197  C14 Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation. <i>The Journal of Cell Biology</i> , 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, <i>Experimental Cell Research</i> , 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. <i>Cell &amp; Tissue Research</i> , 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, <i>Cell</i> , 61:1329-1337 (1990)  Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.	1		PILOSEBACEOUS APPARATUS TO CONTROLLED INJURY, The Journal of Investigative
Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into Complex Programs of Differentiation, <i>The Journal of Cell Biology</i> , 109:2295-2312 (1989)  C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, <i>Experimental Cell Research</i> , 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. <i>Cell &amp; Tissue Research</i> , 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, <i>Cell</i> , 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.			
Complex Programs of Differentiation. <i>The Journal of Cell Biology</i> . 109:2295-2312 (1989)  Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts. <i>Experimental Cell Research</i> . 194: 218-227 (1991)  Limat ea al., Outer root sheath (ORS) cells organize in epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. <i>Cell &amp; Tissue Research</i> , 275:169-176 (1994)  Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, <i>Cell</i> , 61:1329-1337 (1990)  Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C13	Montagna et al., The Structure and Function of Skin 172-258, (Academic Press New York, NY 197-
C15 Limat ea al., Restoration of the Epidermal Phenotype by Follicular Outer Root Sheath Cells in Recombinant Culture with Dermal Fibroblasts, Experimental Cell Research, 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  C17 Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C14	Coulombe et al., Expression of Keratin K14 in the Epidermis and Hair Follicle: Insights into
Recombinant Culture with Dermal Fibroblasts. Experimental Cell Research. 194: 218-227 (1991)  C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.			
C16 Limat ea al., Outer root sheath (ORS) cells organize into epidermoid cyst-like spheroids when cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. Cell & Tissue Research, 275:169-176 (1994)  Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C15	
cultured inside Matrigel: a light-microscipic and immunohistological comparison between human ORS cells and interfollicular keratinocytes. <i>Cell &amp; Tissue Research</i> , 275:169-176 (1994)  Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, <i>Cell</i> , 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.			
ORS cells and interfollicular keratinocytes, Cell & Tissue Research, 275:169-176 (1994)  Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.		C16	
Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implication for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, Cell, 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.			cultured inside Matrigel: a light-microscipic and immunohistological comparison between human
for Follicular Stem Cells, Hair Cycle, and Skin Carcinogenesis, <i>Cell</i> , 61:1329-1337 (1990)  C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.			
C18 Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa.		<b>y</b> ^17	Cotsarelis et al., Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit: Implications
		/	
		C18	Kobayashi, et al., Segregation of keratinocyte colony-forming cells in the bulge of the rat vibrissa,
	<b>*</b>		

Express Mail Label No.: EK674 2461US Date of Deposit: July 10, 2000

E	$\mathcal{L}$		OTHER NON PATENT LITERATURE DOCUMENTS
200		Cite No.	Name of Author, Litle (when appropriate), Publication, Volume, Page(8), Date, Etc.
A	<b>V</b>	C19	Yang et al., Upper Human Hair Follicle Contains a Subpopulation of Keratinocytes with Superior In Vitro Proliferative Potential, The Journal of Investigative Dermatology, 101(5):652-659 (1993)
	<b>*</b>	C20	Rochat, et al., Location of Stem Cells of Human Hair Follicles by Clonal Analysis, Cell, 76:1063-1073 (1994)
		C21	Moll et al., Proliferative Potential of Different Keratinocytes of Plucked Human Hair Follieles, <i>The Journal of Investigative Dermatology</i> , 105:14-21 (1995)
		C22	Weterings, et al., A method for culturing human hair follicle cells, <i>British Journal of Dermatology</i> , 104:1-5 (1981)
		C23	Limat and Noser, Serial Cultivation of Single Keratinocytes from the Outer Root Sheath of Human Scalp Hair Follicles, <i>The Journal of Investigative Dermatology</i> , 87:485-488 (1986)
	1.	C24	Imcke, et al., Growth of human hair follicle keratinocytes in vitro, <i>Journal of the American Academ of Dermatology</i> , 17:779-786 (1983)
		C25	Limat et al., Post-Mitotic Human Dermal Fibroblasts Efficiently Support the Growth of Human Follicular Keratinocytes. <i>The Journal of Investigative Dermatology</i> , 92:758-762 (1989)
		C26	Stark, et al., Keratins of the human hair follicle: "Hyperproliferative" keratins consistently expresse in outer root sheath cells in vivo and in vitro. <i>Differentiation</i> , 35:236-248 (1987)
		C27	Limat et al., Experimental Modulation of the Differentiated Phenotype of Keratinocytes from Epidermis and Hair Follicle Outer Root Sheath and Matrix Cells, <i>Annals of The New York Academy of Sciences</i> , 642:125-147 (1991)
		C28	Lenoir et al., Outer Root Sheath Cells of Human Hair Follicle Are Able to Regenerate a Fully Differentiated Epidermis in Fitro, Developmental Biology, 130:610-620 (1988)
		(*29	Limat et al., FORMATION OF A REGULAR NEO-EPIDERMIS BY CULTRUED HUMAN OUTER ROOT SHEATH CELLS GRAFTED ON NUDE MICE, <i>Transplantation</i> , 59:1032-1038 (1995)
	1.	C30	O'Connor et al., GRAFTING OF BURNS WITH CULTURED EPITHELIUM PREPARED FROM AUTOLOGOUS EPIDERMAL CELLS, <i>The Lancet</i> , 1:75-78 (1981)
		C31	Compton, et al., Skin Regenerated from Cultured Epithelial Autografts on Full-Thickness Burn Wounds from 6 Days to 5 Years after Grafting, <i>Laboratory Investigation</i> , 60:600-612 (1989)
		C32	Carter et al., Treatment of junctional epidermolysis bullosa with epidermal autografts, <i>Journal of th American Academy of Dermatology</i> , 17:246-250 (1987)
	-	C33	Dean et al., The Use of Cultured Epithelial Autograft in a Patient with Idiopathic Pyoderma Gangrenosum, <i>Annals of Plastic Surgery</i> , 26:194-195 (1991)
		C34	Limova and Mauro, Treatment of Pyoderma Gangrenosum with Cultured Keratinocyte Autografts, <i>The Journal of Dermatologic Surgery and Oncology</i> , 20:833-836 (1994)
,		C35	Gallico et al., Cultured Epithelial Autografts for Giant Congenital Nevi, <i>Plastic and Reconstructive Surgery</i> , 84:1-9 (1989)
A	V	C36	Higgins et al., use of two stage keratinocyte-dermal grafting to treat the separation site in conjoined twins, <i>Journal of the Royal Society of Medicine</i> , 87:108-109 (1994)

\* with this reference is not provided as it was previously cited by or submitted to the office in a prior application. U.S.S.N. 09 358.181, filed July 20, 1999, and relied upon for an earlier filing date under 35 U.S.C. §120 (continuation, continuation-in-part, and divisional applications).

	L				
Examiner Signature	NITE	Date Considered	10/2	21/01	

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPFP 609; Draw line through citation if not in conformance and not considered.

Include copy of this form with next communication to applicant.

TRADOCS:1348565.1(SWK501!.DOC)